

Quantification of training and competition load across a season in an elite Australian Football Club

Journal:	International Journal of Sports Physiology and Performance
Manuscript ID:	IJSPP.2015-0294.R2
Manuscript Type:	Original Investigation
Keywords:	training organisation, team sports, training distribution
	•



1 2	Quantification of training and competition load across a season in an elite Australian Football Club
3	
4	Dean Ritchie ^{1,2} , Will G Hopkins ¹ , Martin Buchheit ¹ , Justin Cordy ² , & Jonathan D Bartlett ^{1,2}
5	
6 7	¹ Institute of Sport Exercise and Active Living, Footscray Park,
8	Ballarat Road,
9	Victoria University,
10	Melbourne,
11	Victoria
12	Australia, 8001
13	
14	² Western Bulldogs AFL Club,
14	Whitten Oval,
16	Barkly Street,
17	West Footscray,
18	Victoria,
19	Melbourne,
20	Australia, 3210
21	
22	
23	Running title: Australian Football loading strategies
24	
25	Submission type: Original Investigation
26	
27	
28	Address for correspondence:
29	Institute of Sport Exercise and Active Living,
30	Footscray Park,
31	Ballarat Road,
32	Victoria University,
33	Melbourne,
34	Victoria
35	Australia, 8001
36	Email: Jon.Bartlett@vu.edu.au
37 38	Telephone: +61 3 99 19 4012
50	
39	Word Count: 3,139
40	Abstract word count: 225
41	Number of tables: 2
42	Number of figures: 1

43 Abstract

44 Purpose: Load monitoring in Australian Football (AF) has
45 been widely adopted, yet team sport periodization strategies are
46 relatively unknown. Here we have aimed to quantify training
47 and competition load across a season in an elite AF team, using
48 rating of perceived exertion (RPE) and GPS. tracking.

Methods: Weekly totals for RPE and GPS loads (including
accelerometer data; Playerload) were obtained for 44 players
across a full season for each training modality and for
competition. General linear mixed models compared mean
weekly load between 3x pre-season and 4x in-season blocks.
Effects were assessed with inferences about magnitudes
standardized with between-player SD.

Results: Total RPE load was most likely greater during pre-56 season, where the majority of load was obtained via skills and 57 conditioning. There was a large reduction in RPE load in the 58 last pre-season block. During in-season, half the total load 59 came from games and the remaining half from training, 60 predominantly skills and upper-body weights. Total distance, 61 high-intensity running, and Playerload showed large to very 62 large reductions from pre-season to in-season, whereas changes 63 in mean speed were trivial across all blocks. All these effects 64 were clear at the 99% level. 65

66 **Conclusions**: These data provide useful information about 67 targeted periods of loading and unloading across different 68 stages of a season. Our study also provides a framework for 69 further investigation of training periodization in AF teams.

70

71 **Key Words**: Training organisation, training distribution, team

72 sports

73 Introduction

Australian Football (AF) is a multicyclical competition 74 75 containing a pre-season phase and an in-season phase, requiring athletes to go through a weekly round of competition, recovery, 76 training and subsequent competition.¹ With AF being an 77 intermittent contact sport it requires a wide range of physical 78 attributes such as muscular strength, speed, power, repeated 79 sprint ability, endurance, acceleration, and sport specific 80 skills.^{2, 3} Indeed, players cover anything between 9.5-17 km 81 total distance and in excess of 3 km high-speed (> 14.4 km/h) 82 distance per game.⁴ As such, AF requires careful planning and 83 84 monitoring of training so as to maintain athlete fitness whilst 85 maximising performance.

86

87 The emergence of training load (TL) monitoring in team sports 88 has exponentially grown owing to the need to monitor 89 individual responses to training. Indeed, the adoption of a 90 coach's own training philosophy that is usually based on years of experience and team needs' demonstrates the requirement 91 92 for daily TL evaluation. The use of global positioning systems 93 (GPS) and accelerometers in team sports is now an important 94 monitoring tool for collecting objective information pertaining 95 drills, sessions and games. For example, in-depth to 96 information on the activity profiles of athletes such as total distance travelled, amount of high-intensity running completed, 97 and average movement speed 6, 7 can all be obtained. In 98 addition, the use of the self-perceived session rating (s-RPE) 99 method, known more as a subjective tool, has proved useful in 100 101 determining the internal load of athletes such that the physiological stress to the external load, can effectively be 102 captured.⁸⁻¹⁰ This approach has now been adopted by a number 103 104 of teams as part of their training monitoring system.⁸

105

The ability to obtain both objective and subjective measures of 106 TL allows for a more effective prescription of training. 107 Training periodization requires the careful manipulation of 108 109 training volume and intensity so as to result in an increase in performance.¹¹ Accordingly, the balance between training 110 stress, competition and recovery is of significant importance so 111 as to protect against underperformance ¹² and increased injury 112 risk ¹³. Recent research in soccer and rugby has quantified aspects of weekly^{14, 15}, monthly^{8, 16, 17} and seasonal⁵ TL. 113 114 Despite recent advancements in AF,¹⁸ whereby TL and training 115 duration is higher during pre-season compared to in-season, 116 117 data are limited such that the training and competition load was 118 only quantified using the s-RPE method. The context in which 119 TL is obtained is important as it will allow coaches to better plan and prescribe training at both a team and individual player 120 121 level. As such, information on the external load (alongside that

- of the perceived load) associated with the training practises in
 AF is required. Moreover, where the majority of literature
 compares pre-season to in-season, it is unknown, within these
 two major training and competition phases how load is
 manipulated.
- 127
- 128 The aim of the current study was to quantify training and 129 competition load of a team of Australian Footballers across
- various stages of a season using both s-RPE and GPS.

131 Methods:

132 Subjects

133 Forty-four full-time professional elite AF athletes (mean \pm SD: age, 24.1 ± 3.8 years; height, 187.7 ± 7.2 cm; body mass, 87.3134 \pm 8.2 kg) from the same Australian Football League (AFL) club 135 136 participated in this single full season study. The participating 137 athletes competed in the AFL and the Victorian Football League (VFL) and each provided written informed consent and 138 139 the research was approved by the institutions human research ethics committee. This team achieved a final ranking on the 140 ladder of 14th out of 18 and won 7 and lost 15 games. In the 141 event that players suffered an injury, defined as pain resulting 142 in modified load, data was excluded from the point of injury to 143 144 the point of full return to training.

145 146 **Design**

TL data were collected over a 41 week period during the 2013-147 2014 season. In order to obtain relevant information on training 148 149 and competition loading strategies the season was divided into 150 distinctive periods. Pre-season was sub-divided into pre-season 1, pre-season 2 (divided by the Christmas break)¹⁹ and pre-151 This latter pre-season period incorporated three 152 season 3. 153 practise games. Subsequently, the competition phase was divided into four periods where in-season 1, 3 and 4 contained 154 155 a similar number of games in each with in-season 2 containing 156 no game (bye weekend). Week 26 (in-season 2) was included 157 as its own separate period as it shows how TL is managed during an in-season period when no game is played. The TL 158 presented in each block represents the average weekly total 159 within the given season block so as to account for differences 160 161 in number of weeks within blocks. Individual training sessions, recovery and extras (i.e. individual skill development) were not 162 included in the analysis. In order to analyse the distribution of 163 TL by mode, training was categorised into skills (AF specific 164 165 training), running (field-based conditioning), upper-body weights (UB weights), lower-body weights (LB weights), 166 167 games and "other" (boxing, cycling, swimming and cross-168 training).

169

170 Methodology

Internal TL data were obtained through the RPE-based method 171 ²⁰ at 10-30 minutes following every field-based and indoor 172 training session and games as well as all strength training and 173 cross training conditioning sessions in the gym. In order to 174 obtain a TL value, the RPE is multiplied by session duration, 175 providing a s-RPE for all training and games.²⁰ For all field-176 based training sessions and games, athletes wore GPS devices 177 (MinimaxX S4, Catapult Innovations, Australia). 178 TL 179 parameters obtained from GPS include total distance (m), high-

intensity running (>14.4 km/h (m)) (HIR),²¹ PlayerLoad ²² 180 181 (where the unit of measurement represents the square root of 182 the sum of the squared instantaneous rate of change in acceleration in the X, Y and Z axes divided by 100), and 183 average movement speed (m/min). Each athlete wore the same 184 185 device across the season which was worn inside a custom made vest supplied by the manufacturer across the upper back 186 187 between the left and right scapula. All devices were activated 30-minutes prior to data collection to allow acquisition of 188 189 satellite signals (>8 satellites). The GPS units have a sampling rate of 10 Hz and accelerometer sampling rate of 100 Hz. The 190 accuracy of GPS units sampling at 10 Hz has been shown 191 recently.²³ Following every training session and game, all GPS 192 and accelerometer derived data were downloaded and analysed 193 194 by a specialist GPS software package (Sprint 5.1.3, Catapult 195 Innovations, Australia). A total of 25900 individual training observations and a total of 932 individual game observations 196 197 were obtained. Substitutes in games (N=2 per game) were 198 excluded from the final analysis. Due to the closed roof of the home stadium for 13 of the 26 AFL games full GPS couldn't be 199 200 monitored. However, PlayerLoad was still able to be collected 201 for all games as this was obtained from the accelerometer. All 202 VFL games (N=21) were monitored with both GPS and 203 PlayerLoad, therefore, increasing GPS game sample to N=34. 204 AFL listed players only were included in the analysis.

205 206

207 Statistical analysis:

208 We developed general linear mixed models that estimated 209 training and game loads of players in their uninjured state by 210 including their injury status (total of 41 injuries) as covariates in the model. Covariates were also included to adjust block 211 212 effects to playing position and number of AFL years of 213 experience. Random effects in the model were specified to 214 allow for different between-player standard deviations between 215 blocks (with an unstructured covariance matrix to allow for 216 correlations between blocks) and different within-player 217 standard deviations between blocks (a different residual variance for each block). Effects were assessed with non-218 219 clinical magnitude-based inferences, using standardisation to 220 define magnitude thresholds (lower or equal to 0.20 trivial, lower or equal to 0.60 small, lower or equal to 1.20 moderate, 221 lower or equal to 2.0 large, lower or equal to 4.0 very large and 222 >4.0 extremely large).²⁵ 223 Uncertainty in each effect was 224 expressed as 90% confidence limits (CL) and as probabilities that the true effect was substantially positive or negative.²⁴ To 225 226 account for an inflation of error associated with a large number 227 of inferences in the current study, effects were declared clear at 228 the 99% level.

229 **Results**:

Total RPE Load was most likely greater in pre-season 1 and 2 230 231 than in-season (Table 1, Figure 1). During pre-season 1 and 2 the majority of load most likely came from skills, "other" and 232 233 running in comparison to pre-season 3 and in-season blocks. In contrast, half of the in-season load came from games with the 234 235 remaining half predominantly from skills training and UB weights (Table 1, Figure 1). LB weights were most likely 236 reduced during in-season as was running and "other" 237 conditioning components. 238

Total distance in training was most likely greater during preseason 1 and 2 compared with in-season. In contrast, total
distance covered in games was most likely greater during inseason compared with games in pre-season 3 (Table 2).

244

239

Similar to total distance, there were likely reductions in HIR in
training during in-season compared to pre-season 1 and 2
whilst there was a likely increase in HIR during in-season 3
compared to in-season 1 and 4. Even though HIR was most
likely lower in games during pre-season 3, there was no change
in HIR during games across in-season blocks (Table 2).

Differences in mean speed were most likely trivial for all preseason and in-season blocks for both training and games (Table 2). In contrast, Playerload was most likely higher in training during pre-season 1 and 2 compared to in-season and likely increased during in-season 3 compared with in-season 1 and 4. Playerload in games during pre-season 3 was most likely lower than games during in-season (Table 2).

258 **Discussion**:

259 The aim of the current study was to quantify training and 260 competition load in AF using a combination of s-RPE and GPS 261 load monitoring across specific blocks of a season. We show 262 that load during pre-season was obtained predominantly from 263 conditioning and skills training whereas in-season load was 264 obtained by competition, skills and upper-body weights. At a 265 global level, this is consistent with existing knowledge, where 266 TL is greater during pre-season, whilst in-season there is a 267 concomitant decrease and increase in training and competition 268 load, respectively.

This study is in agreement with existing literature where pre-269 season TL is greater than in-season TL,^{15, 18, 25} however, we 270 271 provide new information in the way in which external load is obtained during the course of a season. Indeed, field-based 272 273 GPS training load was higher in the pre-season compared with in-season, an effect that is likely due to the specific 274 275 conditioning focus of preparing physically for the in-season competition demands. It is well known that pre-season is a 276 277 crucial period for team sports yet it was unclear as to the 278 proportion of work in terms of conditioning and skills they do 279 in the pre-season. Moreover, during the in-season, 280 approximately 50% of external load was obtained by games, 281 whereas the remaining 50% was obtained by training (Figure 282 2b). In contrast to pre-season load distribution though, this in-283 season training load was actually obtained by more skills training and UB weights (Figure 3), whereas in pre-season the 284 training load consisted of high amounts of skills training and all 285 286 aspects of conditioning. Presumably due to the high-intensity 287 nature and increased load of games (~900 RPE load units per 288 game), the difference in in-season training load and the 289 distribution of training mode (i.e. reduction in lower-body load) 290 was likely served to support the recovery process (see below 291 for further information on lower-body load). Whilst the current 292 study did not examine the within-week loading between games, 293 it can be speculated that the reduction in overall training load 294 from pre-season to in-season would also result in a reduction in 295 training load within week, i.e. between games. This 296 periodization strategy is supported by recent work where high 297 training load between both AFL and Rugby League games (separated by 1-week) impairs sprint capacity and explosive 298 actions typical of intermittent activity ¹² and increases injury 299 risk¹³. Together, these data provide important information for 300 practitioners when considering the overall load and mode of 301 302 training that is prescribed to team sport athletes at varying 303 times within a season.

As noted previously, training distribution transitioned from preseason (predominantly running, skills and "other" conditioning) 306 to in-season (skills and UB weights). LB weights load was also 307 greater during the pre-season compared to in-season. Although 308 there may have been a reduction in the frequency of lower body 309 weight sessions during the in-season, it may also be suggested 310 that this reduction in LB weights load was due to an increase in 311 high-intensity running during competition. However, there was 312 a simultaneous decrease in high-intensity running during 313 training in all in-season periods suggesting the reduction in LB 314 weight load is primarily due to the adoption of a recovery 315 focussed training week. Unfortunately, this study is unable to describe whether this dose of LB weights load is capable of 316 317 maintaining or developing strength. Some evidence suggests up to two weekly sessions of strength based training is required 318 for maintenance of muscular strength,²⁶ however, there is 319 limited evidence as to the required dose for elite AF players. 320 321 Future research should aim to uncover the minimal weekly 322 dose required for AF players to maintain a strength and/or 323 hypertrophic stimulus during the in-season period.

324 Unsurprisingly and consistent with the shift in training focus, 325 field-based weekly TL was similar across all in-season blocks. 326 Due to the 1-game per week schedule in AFL, coaches may be 327 able to plan effective in-season training programmes that facilitate the preparation for and recovery from competition.¹² 328 329 Interestingly though, there were only trivial differences in mean 330 speed for training across the duration of both pre-season and in-331 season. This intensity was a lot lower than that of games, 332 highlighting the magnitude of stimulus that games provide. 333 Indeed, the concept of 'train as you play' is highly impractical in this sense owing to the high game demands and increased 334 injury risk. As such, it may actually demonstrate that coaches 335 336 knowingly prescribe an in-season 'maintenance' dose so as to preserve the physical capacities developed during pre-season³² 337 338 but also to ensure optimal preparedness for competition. 339 Furthermore, it may also relate to the reduction in lower-body 340 weights load, such that, more emphasis is placed on 341 maintaining an aerobic fitness stimulus, resulting in a decreased lower-body weights load. It should also be noted that mean 342 343 speed may be particularly dependent on the coach's philosophy, where drills that develop a particular game style 344 345 may be repeated regularly throughout the season. In keeping with this concept of a coach's philosophy, the increase in 346 training duration during in-season 3 may have been a coach 347 348 driven decision targeted to developing game style. 349 Concomitantly, there was also an increase in training HIR and 350 PlayerLoad during in-season 3; a likely result of the increase in 351 duration. These data demonstrate the challenges associated with 352 training design in team sports and may present important 353 questions for coaches and practitioners when planning training 354 during the competitive stage of the season.

Consistent with previous findings,^{18, 27} we report reductions in 355 356 load obtained during pre-season practise games compared to inseason games. This appears to be a direct result of the reduction 357 (approx. 30%) in game time as total distance, HIR and 358 PlayerLoad were also reduced by ~30% suggesting that if game 359 360 time was standardised between pre-season and in-season 361 games, load would have been similar. It may be speculated that 362 coaches adopt a pre-competition reduction in load so as to 363 protect against injury, such as that shown in rugby league 364 where reductions in load in the pre-season reduce risk of injury and result in greater improvements in physical fitness ²⁸. In 365 366 addition, rules on player rotations are also different during 367 practice games compared to AFL competition such that during 368 competitive AFL games, teams are limited to 3 players rolling on and off the ground for a total of 120 rotations per team per 369 370 game. However, during practice games this is unrestricted, where ~ 6 players rotate at any one time with upwards of a total 371 372 of 160-180 rotations. To this end, both training load compared 373 to pre-season 1 and 2 and game load during pre-season 3 compared to in-season is lower. Collectively, these data suggest 374 375 that training and game load is periodically managed prior to 376 competition, possibly in an attempt to reduce risk of injury.

377 Practical applications:

378 The combination of internal (s-RPE) and external (GPS) load monitoring is important for practitioners in understanding all 379 380 load obtained during the course of a season. Indeed, the integration of both internal:external load measures may be a 381 382 viable and feasible monitoring strategy so as to accurately 383 determine loading at various points in the season. Moreover, 384 load distribution is largely affected according to the time of the 385 season, with pre-season containing the highest amounts of 386 conditioning and skills whilst in-season is characterised by a 387 focus on competition and recovery.

Despite these novel findings, it is acknowledged that this is 388 effectively a case study of one team competing in the AFL. 389 390 The authors recognize that the findings are likely specific to this group of players and the specific style and philosophy of 391 392 the coaching staff. As such, further research is required that 393 depicts a broader overview of the TL, intensity and distribution 394 of training in AF. In addition, the training practices presented 395 in the current study are likely to be different at the individual level. That said, load associated with individual skill 396 397 development sessions and recovery should be examined so as to provide an overview of what additional loading these 398 399 provide to the athletes. Furthermore, information on position 400 and years of experience in the AF system as well as the link 401 between performance and injury would provide greater 402 understanding as to the organisation of training and competition

- 403 load during a season and allow for improved athlete 404 conditioning.
- 405 *Conclusion*:

406 This is the first study to systematically quantify the training periodization strategies across a season in Australian Football 407 using both perceived exertion (RPE) and GPS-derived 408 409 monitoring markers. The data from this study revealed that 410 pre-season contains higher training loads, whereas in-season, 411 there is a shift in load distribution such that ~50% of load is 412 obtained via competition. Combined with 'in house' analyses, this distribution of load may aid practitioners in planning and 413 structuring future training plans, as well as to compare and 414 contrast to other practices in Australian Football. As this is an 415 416 analysis of a single team, the distribution and variation of load across the season may vary between clubs. Future research 417 incorporating other modes of load monitoring as well as 418 examining differences in position, AF years of experience and 419 420 individual responses will help our understanding of changes in 421 various components of fitness in response to load.

422

423 Acknowledgements:

The authors declare there was no financial assistance with the preparation of this manuscript nor was there any financial assistance for data collection, data analysis and data interpretation. The authors acknowledge Johann Bilsborough (Western Bulldogs FC) for his constructive comments on the manuscript.

430

431

432	Dofor	ences:
432	1.	Issurin VB. New horizons for the methodology and
434	1.	physiology of training periodisation.pdf>. Sports Med. 2010;
435		40(3):189-206.
436	2.	Bilsborough JC, Greenway KG, Opar DA, et al. Comparison of
437	۷.	anthropometry, upper-body strength, and lower-body
438		power characteristics in different levels of Australian
439		football players. J Strength Cond Res. 2015; 29(3):826-834.
440	3.	Sullivan C, Bilsborough JC, Cianciosi M, Hocking J, Cordy JT,
441	5.	Coutts AJ. Factors affecting match performance in
442		professional Australian football. Int J Sports Physiol Perform.
443		2014; 9(3):561-566.
444	4.	Aughey R. Increased high intensity activity in elite australian
445		football finals matches. Int J Sports Physiol Perform. 2011;
446		6:367-379.
447	5.	Malone JJ, Di Michele R, Morgans R, Burgess D, Morton JP,
448	5.	Drust B. Seasonal Training Load Quantification in Elite
449		English Premier League Soccer Players. Int J Sports Physiol
450		Perform. 2014; 10(4):489-497.
451	6.	Buchheit M, Racinais S, Bilsborough JC, et al. Monitoring
452	-	fitness, fatigue and running performance during a pre-
453		season training camp in elite football players. J Sci Med
454		Sport. 2013; 16(6):550-555.
455	7.	Wisbey B, Montgomery PG, Pyne DB, Rattray B. Quantifying
456		movement demands of AFL football using GPS tracking. J Sci
457		Med Sport. 2010; 13(5):531-536.
458	8.	Scott TJ, Black CR, Quinn J, Coutts AJ. Validity and reliability
459		of the session-RPE method for quantifying training in
460		Australian football: a comparison of the CR10 and CR100
461		scales. J Strength Cond Res. 2013; 27(1):270-276.
462	9.	Gallo T, Cormack S, Gabbett T, Williams M, Lorenzen C.
463		Characteristics impacting on session rating of perceived
464		exertion training load in Australian footballers. J Sports Sci.
465		2015; 33(5):467-475.
466	10.	Gaudino P, Iaia FM, Strudwick AJ, et al. Factors Influencing
467		Perception of Effort (Session-RPE) During Elite Soccer
468		Training. Int J Sports Physiol Perform. 2015 In Press.
469	11.	Meeusen R, Duclos M, Foster C, et al. Prevention, diagnosis,
470		and treatment of the overtraining syndrome: joint
471		consensus statement of the European College of Sport
472		Science and the American College of Sports Medicine. Med
473		Sci Sports Exerc. 2013; 45(1):186-205.
474	12.	Slattery KM, Wallace LK, Bentley DJ, Coutts AJ. Effect of
475		training load on simulated team sport match performance.
476	4.5	Appl Physiol Nutr Metab. 2012; 37(2):315-322.
477	13.	Murray NB, Gabbett TJ, Chamari K. Effect of different
478		between-match recovery times on the activity profiles and
479		injury rates of national rugby league players. J Strength Cond
480	14	Res. 2014; 28(12):3476-3483.
481 482	14.	Coutts A, Reaburn P, Piva TJ, Murphy A. Changes in selected
482		biochemical, muscular strength, power, and endurance

400		management during deliberate everyophing and tenewing in
483		measures during deliberate overreaching and tapering in
484		rugby league players. Int J Sports Med. 2007; 28(2):116-124.
485	15.	Jeong TS, Reilly T, Morton J, Bae SW, Drust B. Quantification
486		of the physiological loading of one week of "pre-season" and
487		one week of "in-season" training in professional soccer
488		players. J Sports Sci. 2011; 29(11):1161-1166.
489	16.	Impellizzeri FM, Rampinini E, Coutts AJ, Sassi A, Marcora SM.
490		Use of RPE-Based Training Load in Soccer. Medicine &
491		Science in Sports & Exercise. 2004; 36(6):1042-1047.
492	17.	Casamichana D, Castellano J, Calleja-Gonzalez J, San Roman
493		J, Castagna C. Relationship between indicators of training
494		load in soccer players. J Strength Cond Res. 2013; 27(2):369-
495		374.
496	18.	Moreira A, Bilsborough JC, Sullivan CJ, Ciancosi M, Aoki MS,
497		Coutts AJ. The Training Periodization of Professional
498		Australian Football Players During an Entire AFL Season. Int J
499		Sports Physiol Perform. 2014, 10(5): 566-571.
500	19.	Buchheit M, Morgan W, Wallace J, Bode M, Poulos N.
501		Physiological, psychometric, and performance effects of the
502		Christmas break in Australian football. Int J Sports Physiol
503		Perform. 2015; 10(1):120-123.
504	20.	Foster C, Florhaug JA, Franklin J, et al. A new approach to
505		monitoring exercise training. J Strength Cond Res. 2001;
506		15(1):109-115.
507	21.	Coutts AJ, Quinn J, Hocking J, Castagna C, Rampinini E.
508		Match running performance in elite Australian Rules
509		Football. J Sci Med Sport. 2010; 13(5):543-548.
510	22.	Barrett S, Midgley A, Lovell R. PlayerLoad: reliability,
511		convergent validity, and influence of unit position during
512		treadmill running. Int J Sports Physiol Perform. 2014;
513		9(6):945-952.
514	23.	Rampinini E, Alberti G, Fiorenza M, et al. Accuracy of GPS
515		devices for measuring high-intensity running in field-based
516		team sports. Int J Sports Med. 2015; 36(1):49-53.
517	24.	Hopkins WG, Marshall SW, Batterham AM, Hanin J.
518		Progressive statistics for studies in sports medicine and
519		exercise science. Med Sci Sports Exerc. 2009; 41(1):3-13.
520	25.	Rogalski B, Dawson B, Heasman J, Gabbett TJ. Training and
521		game loads and injury risk in elite Australian footballers. J Sci
522		Med Sport. 2013; 16(6):499-503.
523	26.	McMaster DT, Gill N, Cronin J, McGuigan M. The
524		development, retention and decay rates of strength and
525		power in elite rugby union, rugby league and American
526		football: a systematic review. <i>Sports Med.</i> 2013; 43(5):367-
527		384.
528	27.	Colby MJ, Dawson B, Heasman J, Rogalski B, Gabbett TJ.
529		Accelerometer and GPS-derived running loads and injury
530		risk in elite Australian footballers. J Strength Cond Res. 2014;
531		28(8):2244-2252.

- 532 28. Gabbett TJ. Reductions in pre-season training loads reduce
- 533 training injury rates in rugby league players. *Br J Sports Med.*
- 534 2004; 38(6):743-749.

535

Table 1. Quantification of weekly training and game load
throughout each block during the season for total, games, skills,
UB weights, LB weights, other and running load. Standardised
differences are denoted by letters and expressed by effect size.
Data are shown as mean ± SD.

Table 2. Quantification of weekly training and game load
throughout each block during the season for duration, total
distance, high-intensity running, mean speed, and PlayerLoad.
Standardised differences are denoted by letters and expressed
by effect size. Data are shown as mean ± SD.

546 Figure 1. Training distribution expressed by RPE Load per 547 week within block for weekly total load (large bar) and all modes (small bars). Pre-season 1 and pre-season 2; M denotes 548 moderate standardised difference vs in-season 1, 3 and 4; L 549 denotes large standardised difference vs pre-season 3 and in-550 551 season 2. Pre-season 3; S denotes small standardised difference vs in-season 1, 3 and 4. In-season 2; M denotes moderate 552 standardised difference vs in-season 1, 3 and 4. Data are shown 553 554 as mean \pm SD.

Human Kinetics, 1607 N Market St, Champaign, IL 61825

TABLE 1.

Table 1. Quantification of weekly training and game load distribution throughout each block during the season for total, games, skills, UB weights, LB weights, other andrunning.Standardised differences are denoted by letters and expressed by effect size.Data are shown as mean \pm SD.

Block	Total (AU)	Games (AU)	Skills (AU)	UB weights (AU)	LB weights (AU)	Other (AU)	Running (AU)
Pre-Season 1	$2740 \pm 1330^{\ M, \ L}$	-	$600\pm470^{~S,~L}$	$370\pm200^{-S,\ L}$	$390\pm 200^{~S,~M,~L}$	$740 \pm 530^{8, L}$	$640\pm1080^{\text{ M, L}}$
Pre-Season 2	$2680 \pm 710^{\ M,\ L}$	-	$1090\pm490~^L$	$320\pm170^{\ M,\ L}$	$420\pm270^{\rm L}$	$610\pm500^{\rm \ L}$	$220\pm240~^{\rm S}$
Pre-Season 3	$1570 \pm 540^{-S,\ M}$	$570\pm240~^{\rm L}$	520 ± 340	$150\pm140^{~M,L}$	210 ± 90^{8}	160 ± 180	110 ± 170
In-Season 1	1950 ± 600	940 ± 180	480 ± 220	$280\pm130^{\ S,M}$	150 ± 90	150 ± 250	40 ± 170
In-Season 2	$1460\pm340^{\rm \ M}$		$410\pm140\ ^{S}$	$420\pm180^{~S,M}$	140 ± 40	140 ± 310	$270\pm210\ ^{\rm S}$
In-Season 3	2130 ± 520 ^s	970 ± 180^{-8}	580 ± 250	$370\pm180\ ^{\mathrm{S}}$	160 ± 80	130 ± 170	50 ± 120
In-Season 4	1870 ± 580	980 ± 190	470 ± 180	330 ± 130	160 ± 100	90 ± 200	50 ± 150

Superscripts indicate small (S), moderate (M), large (L) and very large (V) differences (clear at the 99% level) as follows.

Total:

Pre-season 1 and Pre-season 2; M vs in-season 1, in-season 3 and in-season 4. L vs pre-season 3 and in-season 2.

Pre-season 3; S vs in-season 1 and in-season 4. M vs in-season 3.

In-season 2; M vs in-season 1, in-season 3 and in-season 4.

In-season 3; S vs in-season 1 and in-season 4

Game:

L vs all in-season blocks and S vs in-season 1.

Skills:

Pre-season 1; S vs in-season 2 and in-season 4. L vs pre-season 2.

Pre-season 2; L vs all in-season blocks

In-season 2; S vs in-season 3

UB weights:

Pre-season 1; S vs pre-season 2, in-season 1, in-season 2 and in-season 4. L vs pre-season 3.

Pre-season 2; M vs pre-season 3 and in-season 3. L vs in-season 2.

Pre-season 3; M vs in-season 1 and in-season 4. L vs in-season 2 and in-season 3.

In-season 1; S vs in-season 3 and in-season 4. M vs in-season 2.

In-season 2; S vs in-season 3. M vs in-season 4.

In-season 3; S vs in-season 4.

LB weights:

Preseason 1; S vs pre-season 2, M vs pre-season 3 and L vs all in-season blocks.

Pre-season 2; L vs pre-season 3 and all in-season blocks

Pre-season 3; S vs all in-season blocks



Other:

Pre-season 1 and 2; S vs pre-season 2 and L vs pre-season 3 and all in-season blocks.

Running:

Pre-season 1; M vs pre-season 2, pre-season 3 and in-season 2. L vs in-season 1, in-season 3 and in-season 4.

Pre-season 2; S vs pre-season 3, in-season 1, in-season 3 and in-season 4.

In-season 2; S vs pre-season 3, in-season 1, in-season 3 and in-season 4.

Table 2. Quantification of weekly training and game load throughout each block during the season for duration, total distance, high-intensity running, mean speed,PlayerLoad and maximal accelerations. Standardised differences are denoted by letters and expressed by effect size. Data are shown as mean \pm SD.

	Duration (min)		Total Distance (m)		HIR (m)		Mean Speed (m/min)		PlayerLoad (AU)	
Block	Training	Game	Training	Game	Training	Game	Training	Game	Training	Game
Pre-Season 1	$199\pm76~^{\rm L}$	-	$20000 \pm 8200^{\text{L}}$	-	6680 ± 3540 ^{LV}	-	99 ± 201	-	$1910\pm770~^{\rm L}$	-
Pre-Season 2	$209\pm72~^{\rm L}$	-	21400 ± 7300 ^L	-	$6350 \pm 2490 \ ^{\rm LV}$	-	101 ± 152	-	$2060\pm720\ ^{\rm L}$	-
Pre-Season 3	103 ± 49	$69\pm21~^{\rm V}$	10200 ± 5600	9900 ± 3000 ^L	2630 ± 2120	$2550\pm840\ ^{M}$	98 ± 149	142 ± 73	1000 ± 500	1010 ± 290^{L}
In-Season 1	112 ± 41	100 ± 13	9900 ± 3800	13300 ± 1700	2440 ± 1120	3140 ± 820	87 ± 102	132 ± 80	980 ± 380	1310 ± 190
In-Season 2	117 ± 24	-	10500 ± 2500	6	2850 ± 1050	-	88 ± 79	-	970 ± 210	-
In-Season 3	126 ± 52 ^s	101 ± 13	11800 ± 4400 ^s	13400 ± 1500	2970 ± 1400 ^s	3270 ± 670	93 ± 128	132 ± 74	1130 ± 430 ^s	1320 ± 190
In-Season 4	111 ± 38	102 ± 14	10400 ± 3300	13500 ± 1700	2430 ± 900	3330 ± 810	93 ± 78	133 ± 57	990 ± 320	1320 ± 200

Superscripts indicate small (S), moderate (M), large (L) and very large (V) differences (clear at the 99% level) as follows.

Training Duration: L vs pre-season 3 and all in-season blocks and S vs pre-season 3.

Game Duration: V vs all in-season blocks.


Training Total Distance: L vs pre-season 3 and all in-season blocks and S vs pre-season 3 and in-season 1.

Game Total Distance: L vs all in-season blocks.

Training High-Intensity Running: LV vs pre-season 3 and all in-season blocks and S vs in-season 1 and in-season 4. Game High-Intensity Running: M vs all in-season blocks.

Training Player Load: L vs pre-season 3 and all in-season periods and S vs in-season 1, in-season 2 and in-season 4. Game Player Load: L vs all in-season blocks.

FIGURE 1



244x118mm (150 x 150 DPI)

