



THE QUADRANT OF DOOM AND HAMSTRING INJURIES: SEXY BUT TOO EASY?

FEATURE / BUCHHEIT M, AVRILLON S, SIMPSON B.M, LACOME M & GUILHEM G

Introduction

The different factors relating to hamstring injury risk have been well reviewed. They include among others; age, previous injuries, ethnicity, strength and strength imbalances, flexibility, muscle architecture, anatomy, training/competitive load (often high-speed running) and fatigue⁽¹⁻³⁾. Recently, there has been a growing emphasis on two of those factors, namely hamstring strength and fascicle length⁽⁴⁻⁶⁾. This is related to the fact that these two muscle properties are modifiable factors strongly related to the capacity of the muscle to withstand repeated eccentric contractions during potentially harmful actions such as sprinting. In fact, it has been suggested that players with weak knee flexor eccentric strength (as measured using a Nordbord, Vald Performance, Brisbane, Australia) and short biceps femoris long head (BF_{lh}) fascicle length may be at much greater risk of injury than players with strong knee flexors and long fascicle length⁽⁶⁾. This has led some authors to present the data in the form of a "quadrant of doom"

⁽⁷⁾, where the overall risk of an individual sustaining a hamstring injury is shown graphically, while plotted as a function of both hamstring strength and fascicle length. It is therefore understood that athletes should escape from the lower left quadrant (high risk), and enter the top right panel of the graph (lower risk), likely via eccentric-biased training⁽⁷⁾. The idea behind the quadrant is evidence-based and sensible⁽⁸⁾, and the highly practical aspect of those strength and structural measures make the approach very appealing for practitioners. Nonetheless, we wished to comment on two important and still overlooked methodological aspects that deserve more attention to make the most of the utilisation of the quadrant: 1) the possible impact of body mass (BM) on Nordbord performance⁽⁹⁻¹¹⁾ 2) the current limitations of the muscle architecture measurements inferred from static ultrasound images and 3) possible differences in individual muscle properties and their relationships with hamstring ability to withstand active lengthening^(12, 13).

The impact of Body Mass on Nordbord performance

The need to consider players' BM when it comes to assessing Nordbord performance is straightforward for most practitioners (Figure 1), and is not a new finding in the scientific literature either⁽⁹⁻¹¹⁾. At least three independent studies have now reported moderate-to-large relationships between Nordbord performance and BM, and have shown -although correlations don't imply causality- that Nordbord performance likely increases consistently by 3⁽¹¹⁾ to 4^(9, 10) N per kg of BM. This is not surprising, for at least two reasons: 1) for most neuromuscular-related types of measures, including hamstring strength⁽¹⁴⁾, muscle mass is generally beneficial for performance⁽¹⁵⁾ and 2) because of the upper body inclination when leaning forward during the Nordic exercise, heavier and/or taller players with a longer lower-leg lever (distance from knee joint axis of rotation to the ankle strap) may apply higher levels of force to the dynamometers. Greater Nordbord performance in heavier

players may be in turn interpreted as a greater eccentric knee flexor strength, which may be independent (at least partially) of their true strength. It is however important to note that the beneficial effect of a greater BM on Nordbord performance may only be apparent for the players that are strong enough to perform the exercise in a controlled manner, since more load added to the chest in athletes with weak knee flexors eccentric strength will likely only make them fall faster, with no effect on Nordbord performance. However, and while we agree that the suggested normalising procedure⁽¹⁰⁾ still lacks prospective evidence, until a new solution is provided, scaling Nordbord performance for BM remains the most practical way to account for this likely confounding factor. While we also agree that the value of the reductions reported in those three studies (3⁽¹¹⁾ to 4^(9, 10) N per kg of BM) may not be as steep in more homogenous/different players groups (unpublished data from⁽⁶⁾), we still believe that this relationship should be tested and then accounted for if present (irrespective of its magnitude). Surprisingly, despite this evidence, most researchers have continued to report absolute strength values (N) in their studies^(1, 4-6). They have also used a unique absolute eccentric strength threshold value to identify players with increased hamstring injury risk (i.e., 265 N)⁽¹⁶⁾ or to design the quadrant (i.e., 337 N, in Figures 2 and 3 in ref⁽¹⁾), without taking their own BM into consideration; therefore, this procedure remains prone to approximations. It is also important to note that simply dividing eccentric strength by units of BM (i.e., N/kg)^(9, 16) is unlikely optimal either. The levels of correlation (and slope magnitudes) reported in players of different ages and sports^(9, 10) suggest that the relationship between eccentric knee flexor strength and body size (and likely muscle architecture, see next section) is complex, and likely be specific to

the group of players considered (i.e., group-based allometric scaling parameters⁽¹⁷⁾). Overall, this data suggests that BM should not be overlooked when monitoring Nordbord performance, which may limit, at least in theory, the usefulness of the “quadrant of doom” as currently presented^(1, 7). Further studies are nevertheless required to confirm whether BM-adjusted strength values improve injury risk prediction in elite soccer players.

To further illustrate our point, we used data recently collected in young elite footballers⁽¹⁸⁾ and reproduced a typical “quadrant of doom”⁽¹⁾ using the suggested absolute strength threshold (i.e., 337 N Figure 2, left panel). While we agree that the value of such a cut-off is often sample-dependent, the point we are trying to make here is likely valid irrespective of the actual value chosen. Following this initial reasoning, player #19 (56 kg, knee-flexor strength: 316 N, BFLh fascicle length: 8.1 cm, bottom left quadrant) was reported to have a higher risk of injury than player #2 (73 kg, 384.5 N; 7.5 cm, lower right quadrant). However, after adjusting players’ knee flexors strength to their BM⁽¹⁷⁾, completely different figures were apparent, with risk profiles being drastically different, i.e., player #19’s relative strength (compared with body-mass expected performance): +27.1%, player #2: +16.6%. While player #2 remained in the same quadrant, player #19 moved into the lower right quadrant, which likely signified lower injury risk. With this particular player (#19), for example, practitioners (ourselves in this case!) would clearly face a dilemma when assessing his injury risk.

Hamstring muscle properties and their relative susceptibility to injury. The second methodological point that we wished to comment on is related to the other axis (i.e., fascicle length). There are several points that deserve consideration:



Overall, this data suggests that BM should not be overlooked when monitoring Nordbord performance, which may limit, at least in theory, the usefulness of the ‘quadrant of doom’ as currently presented.”



a. In fact, players' anthropometrical profile not only impacts on muscle mass, but also on muscle geometry and size, since fascicle length, muscle thickness and pennation are interconnected factors. For example, the fascicle length of a fusiform muscle such as the semitendinosus is likely directly related to the length of the femur and in turn, to the player's size⁽¹⁹⁾. This suggests that taller players are likely to present with longer fascicle length, which may have nothing to do with the muscle's ability to withstand active lengthening per se. Therefore, as for strength measures, body size likely confounds the relationship between fascicle length and players' actual injury risk. Normalizing fascicle length for muscle length^(6, 19) may therefore constitute, at least in theory, a first improvement to the only use of absolute fascicle length⁽¹⁾. In contrast to this reasoning however, normalized measures of fascicle length were pretty similarly related to injury rates in the unique study to date in soccer players⁽⁶⁾. It is however worth noting that overall group-based results may not always apply to extreme case scenarios (Figure 1); therefore, additional studies in players of varying sizes are necessary to clarify this point.

b. Importantly, fascicle length measurements of pennate muscles such as the BFH with a 4.7-cm probe also require a substantial extrapolation ($\approx 60\%$ of the entire length for

a 11.8 cm-long fascicle⁽²⁰⁾), which can lead to a $3^{(20)}$ to $5^{(21)}$ % error that is unfortunately greater than the smallest important effect (i.e., $0.2 \times$ between-subject SD⁽²²⁾), estimated to be around 2% in our population⁽¹⁸⁾). This poor signal/noise ratio shows the limitation of using a single fascicle length measurement on the basis of a single B-mode image to draw the quadrant. To improve precision and in turn, confidence in their assessment, practitioners may therefore need to use i) repeated measures that can decrease the noise by a factor of \sqrt{n} ⁽²³⁾ or ii) alternate muscle architecture measurements using MRI⁽²⁴⁾, diffusion tensor imaging⁽²⁵⁾ or extended field of view (EFOV) measures⁽¹⁸⁾ for example. This latter mode uses an algorithm that fits series of images, allowing scanning of entire fascicles within one continuous scan. This technique may therefore enable practitioners to avoid any extrapolation of non-visible parts of the muscle and provides improved measurement accuracy^(21, 26). Using a scan that follows fascicle orientations along their path (non-linear EFOV) can further account for fascicle curvature and improve the imaging of the fascicle, particularly in the distal regions, resulting in higher reliability compared to single B-mode images or linear EFOV (Pimenta et al. 2018). Interestingly, this method revealed reasonable increases in BFH fascicle length ($\sim +0.5$ cm i.e. $+5\%$) in elite football

players after 6 weeks of eccentric-biased hamstring training⁽¹⁸⁾.

c. Despite the evidence showing the relationship between injuries and fascicle length measured in a relaxed state⁽¹⁾ (i.e., in a resting position as we have also done, Figure 2) it is worth noting that such an assessment is unlikely to accurately represent an individual muscle's ability to withstand an active lengthening. This is particularly true for a muscle group with such a complex and heterogeneous architecture as the hamstring^(19, 27), which may likely involve fascicle rotations during contractions⁽²⁸⁾. In addition, while it is true that fascicle lengthening is related to functional alterations induced by exercise⁽²⁹⁾, the elastic properties of tendinous tissue may mitigate the extent of fascicle strain^(30, 31). Moreover, fascicle length was considered for BFH only, whilst knee flexor strength measured at the joint level reflects the contribution of all synergists and antagonist muscles. In fact, for the same joint motion, the semitendinosus likely displays less relative strain than the other hamstrings probably owing to a greater length, longer fascicles and, possibly, a longer tendon⁽¹⁹⁾. Using kinematics and ground reaction force data integrated with a three-dimensional musculoskeletal computer model, Schache et al.⁽¹³⁾




Figure 2. In-season values of elite U19 soccer players (17.5 ± 0.7 yrs, 175.7 ± 5.0 cm and 64.7 ± 4.9 kg, training 10 hrs a week) for biceps femoris long head (upper panels, BFH, y-axis) and semimembranosus (lower panels, y-axis) fascicle length and eccentric knee flexor strength (Nordbord performance, x-axis) expressed in absolute (left panels) and relative to body mass (right panels). The eccentric knee-flexor strength testing was performed as previously described⁽¹⁰⁾. As between-leg differences were beyond the scope of the current study, the average strength of both legs was used for analysis⁽¹⁰⁾. The data relative to body mass are expressed as the % difference vs. body-mass expected value using the following equation: eccentric strength (N) = $4 \times \text{BM (kg)} + 26.1$ ⁽¹⁰⁾. While we agree that the adjustment of Nordbord performance for BM may be optimal using population-specific equations⁽¹⁰⁾, we chose to use this generic equation since this is what most practitioners would use initially, before getting their own equation. We also believe that using a group-specific equation would not change the main message of the present example. Muscle fascicles were imaged using a 42-mm linear probe (2–10 MHz, SL10-2, Supersonic Imagine, Aix-en-Provence, France) coupled with an ultrasound scanner (Aixplorer V11, Supersonic Imagine, Aix-en-Provence, France)⁽¹⁸⁾. Given that the field-of-view of the probe was too narrow to image an entire fascicle, we used an inbuilt panoramic mode of the ultrasound device. This mode uses an algorithm that fits series of images, allowing scanning of entire fascicles within one continuous scan. This technique enabled us to avoid any extrapolation of non-visible parts of the muscle and improved measurement accuracy⁽²¹⁾. We used this scan to measure the length of two fascicles per muscle. The two values were then averaged to obtain a representative value for the whole muscle. Reliability assessment in our laboratory ($n = 12$, test-retest within 24h) showed small and trivial day-to-day variations in BFH length (typical error: 0.38 ± 0.15 cm, $4.9 \pm 2.0\%$). Dotted lines for strength are based on recommend thresholds⁽¹⁾. Because of the difference in methods used to measure BFH length in comparison with the literature, the dotted line was based on the median value of the current group (i.e., 8.7 cm).




Figure 1. Relationship (r with 90% confidence intervals) between eccentric knee flexor strength and body mass (BM) in six teams. TEE: typical error of the estimate, with 90% confidence intervals. The different lines represent threshold for slightly, moderately and largely lower/greater than BM-expected strength, based on Cohen's effect size principle. For individuals, difference vs. group mean are generally considered as substantial when the probabilities are $\geq 75\%$, which occurs when the difference is greater than the sum of the smallest worthwhile difference (SWD, = TEE/5) and the typical error of measurement (TE, from reliability studies). Reproduced with permission from (10). While player B (106 kg) demonstrates one of the highest level of absolute strength (451 N), in comparison with his BM-expected strength (451 N), his relative strength (+10%) is actually lower than that of player A (75kg, 429 - 326 N = +32%).

suggested that during sprinting, "the BFlh exhibited the largest peak strain, the semitendinosus displayed the greatest lengthening velocity, and the semimembranosus produced the highest peak force, absorbed and generated the most power, and performed the largest amount of positive and negative work". These findings highlight nicely the distinct contributions of each muscle head to lower-limb kinetics during running. Recent studies have also demonstrated that the distribution of force between the heads of the different thigh muscles is often highly variable between individuals⁽³²⁾. The direct consequence of this is that the relative load sustainable by each hamstring head for the same (measured) knee flexor strength may also vary between players⁽³³⁾. Therefore, since the relationship between fascicle length, muscle strength and strain during active lengthening is probably muscle head- and player-dependent, the use of a single measure (i.e., fascicle length) on a single muscle (e.g., BFlh) to assess injury risks remains imperfect, even though a majority of injuries incurred during high speed running occur within the BFlh. Although the single-muscle approach is appealing and particularly adapted to on-field conditions encountered in elite sport, further investigations are required to better understand the individual relationships between the properties of muscle-

tendon unit and force-generating capacity of each hamstring muscle. The consideration of these biomechanical features may in turn contribute to a better evaluation of the injury risk of each individual muscle and a greater individualization of prevention programs⁽¹²⁾. In fact, assessing the properties of each hamstring muscle should give us more information and may be, in turn, more useful in preventing injuries.

d. In practice, the aforementioned differences in muscle properties within the hamstring group are another important limitation to the use of the quadrant as currently presented (Figure 2). In fact, because of the variations in length and structure between the hamstring muscles, a player's position within the quadrant may vary as a function of the muscle considered. For example, player #12 moves from the bottom left (higher risk) to the upper left quadrant (lower risk) when considering the semimembranosus or the BFlh, respectively. Player #10 remains in the same quadrant but moves from a position close to the lower quadrant (semimembranosus) to the highest y-axis position of the group (BFlh). As for absolute vs. relative strength, the fact that players may move from a quadrant to another in relation to the muscle considered represents an important challenge for practitioners seeking a robust means of assessing injury risk.

Additional considerations. Lastly, as the "quadrant of doom" is only a two-dimensional representation of hamstring injury risk, other important risk factors such as age and previous injury history⁽⁶⁾ can't be integrated into the 'picture'. This is another important limitation of the "quadrant of doom" as currently presented.

Conclusion. To conclude, while our intention is definitely not to discard the proposed approach (quadrant) and on-field methodology (easy and quick Nordbord testing and echography measures) that are particularly relevant for practitioners, we wished to highlight some of the limitations that may need to be considered for a better understanding of players' potential injury risk. The example presented in the present paper (Figure 2) suggests the need for considering at least (i) BM when assessing knee-flexors eccentric strength using a Nordbord⁽⁹⁻¹¹⁾ and (ii) individual muscle-tendon properties when estimating hamstring ability to withstand active lengthening. More specifically, we believe that the effect of these two intrinsic factors should not be overlooked when assessing injury risk using a quadrant⁽¹⁾. Understanding these limitations should help practitioners to make better decisions and implement targeted injury prevention interventions for high risk players.

Key points

- The idea behind the “quadrant of doom” is evidence-based and sensible, and the highly practical aspect of those muscle strength and architecture measures make the approach very appealing for practitioners.
- However, the importance of body mass should not be overlooked when monitoring Nordbord performance, which may limit the relevance of the “quadrant of doom” as currently provided with absolute strength values.
- Similarly, since body size may also directly affect muscle length, it is intuitive to normalize the fascicles length used in the “quadrant of doom” for their relative muscle length.
- The measurement of fascicle length with the 2D static image technique likely overestimates fascicle length when compared to extended field of view techniques, thereby affecting the subsequent muscle function and injury risk estimates.
- Since the relationship between fascicle length, muscle strength and strain during active lengthening is probably muscle head- and player-dependent, the use of a single measure (i.e., fascicle length) on a single muscle (e.g., biceps femoris long head, BF_{lh}) to assess the overall injury risk of the hamstring group remains questionable.
- Considering that the “quadrant of doom” is a two-dimensional representation of hamstring injury risk factors, other important risk factors such as age and previous injury history can't be integrated into the 'picture'; this can bias the risk evaluation.
- Although sound in theory, whether the aforementioned theoretical arguments substantially improve the prognostic value of the “quadrant of doom” when it comes to predicting injuries remains to be investigated with real data. More research is still warranted to both improve 1) our understanding and use of Nordbord performance in relation to body mass, and 2) the prognostic value of isolated muscle properties in relation to the overall hamstring group.

1. Bourne, M.N., et al., An Evidence-Based Framework for Strengthening Exercises to Prevent Hamstring Injury. *Sports Med*, 2018. 48(2): p. 251-267.
2. Duhig, S., et al., Effect of high-speed running on hamstring strain injury risk. *Br J Sports Med*, 2016.
3. Opar, D.A., M.D. Williams, and A.J. Shield. Hamstring Strain Injuries: Factors that Lead to Injury and Re-Injury. *Sports Med*, 2012. 42(3): p. 209-26.
4. Bourne, M.N., et al., Impact of the Nordic hamstring and hip extension exercises on hamstring architecture and morphology: implications for injury prevention. *Br J Sports Med*, 2017. 51(5): p. 469-477.
5. Presland, J.D., et al., The effect of Nordic hamstring exercise training volume on biceps femoris long head architectural adaptation. *Scand J Med Sci Sports*, 2018.
6. Timmins, R.G., et al., Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. *Br J Sports Med*, 2016. 50(24): p. 1524-1535.
7. Bourne, M.N., et al., Infographic. Impact of the Nordic hamstring and hip extension exercises on hamstring architecture and morphology: implications for injury prevention. *Br J Sports Med*, 2018.
8. Shield, A.J. and M.N. Bourne. Hamstring Injury Prevention Practices in Elite Sport: Evidence for Eccentric Strength vs. Lumbo-Pelvic Training. *Sports Med*, 2018. 48(3): p. 513-524.
9. Roe, M., et al., Eccentric knee flexor strength profiles of 341 elite male academy and senior Gaelic football players: Do body mass and previous hamstring injury impact performance? *Phys Ther Sport*, 2018.
10. Buchheit, M., et al., The Effect of Body Mass on Eccentric Knee-Flexor Strength Assessed With an Instrumented Nordic Hamstring Device (Nordbord) in Football Players. *Int J Sports Physiol Perform*, 2016. 11(6): p. 721-726.
11. Vald Performance, Premier league & championship hamstring strength report - 2016-17 season. Internal document. 7p. 2017.
12. Erteit, T. and T. Gronwald, M. biceps femoris - A wolf in sheep's clothing: The downside of a lower limb injury prevention training. *Med Hypotheses*, 2017. 109: p. 119-125.
13. Schache, A.G., et al., Mechanics of the human hamstring muscles during sprinting. *Med Sci Sports Exerc*, 2012. 44(4): p. 647-58.
14. Zakas, A., et al., Peak torque of quadriceps and hamstring muscles in basketball and soccer players of different divisions. *J Sports Med Phys Fitness*, 1995. 35(3): p. 199-205.
15. Nedeljkovic, A., et al., Tests of muscle power output: the role of body size. *Int J Sports Med*, 2009. 30(2): p. 100-6.
16. Opar, D.A., et al., Eccentric hamstring strength and hamstring injury risk in Australian footballers. *Med Sci Sports Exerc*, 2015. 47(4): p. 857-65.
17. Jaric, S., D. Mirkov, and G. Markovic, Normalizing physical performance tests for body size: a proposal for standardization. *J Strength Cond Res*, 2005. 19(2): p. 467-74.
18. Lacombe, M., et al., Knee-flexor eccentric training: Does training volume really matter? *Int J Sports Physiol Perform*, 2019: p. Apr 29:1-27. doi: 10.1123/ijsp.2018-0947. [Epub ahead of print].
19. Kellis, E., Intra- and Inter-Muscular Variations in Hamstring Architecture and Mechanics and Their Implications for Injury: A Narrative Review. *Sports Med*, 2018. 48(10): p. 2271-2283.
20. Timmins, R.G., et al., Biceps femoris long head architecture: a reliability and retrospective injury study. *Med Sci Sports Exerc*, 2015. 47(5): p. 905-13.
21. Adkins, A.N., P.W. Franks, and W.M. Murray, Demonstration of extended field-of-view ultrasound's potential to increase the pool of muscles for which in vivo fascicle length is measurable. *J Biomech*, 2017. 63: p. 179-185.
22. Buchheit, M., Magnitudes matter more than Beetroot Juice. *Sport Performance & Science Reports*, 2018. Janv(15): p. vi.
23. Buchheit, M., The Numbers Will Love You Back in Return-Promise. *Int J Sports Physiol Perform*, 2016. 11(4): p. 551-4.
24. Oudeman, J., et al., A novel diffusion-tensor MRI approach for skeletal muscle fascicle length measurements. *Physiol Rep*, 2016. 4(24).
25. Bolsterlee, B., et al., Comparison of measurements of medial gastrocnemius architectural parameters from ultrasound and diffusion tensor images. *J Biomech*, 2015. 48(6): p. 1133-40.
26. Franchi, M.V., et al., Muscle Architecture Assessment: Strengths, Shortcomings and New Frontiers of in Vivo Imaging Techniques. *Ultrasound Med Biol*, 2018. 44(12): p. 2492-2504.
27. Kellis, E., et al., Muscle architecture variations along the human semitendinosus and biceps femoris (long head) length. *J Electromyogr Kinesiol*, 2010. 20(6): p. 1237-43.
28. Azizi, E. and T.J. Roberts, Geared up to stretch: pennate muscle behavior during active lengthening. *J Exp Biol*, 2014. 217(Pt 3): p. 376-81.
29. Guilhem, G., et al., Muscle force loss and soreness subsequent to maximal eccentric contractions depend on the amount of fascicle strain in vivo. *Acta Physiol (Oxf)*, 2016. 217(2): p. 152-63.
30. Hicks, K.M., et al., Gender differences in fascicular lengthening during eccentric contractions: the role of the patella tendon stiffness. *Acta Physiol (Oxf)*, 2013. 209(3): p. 235-44.
31. Konow, N., et al., Spring or string: does tendon elastic action influence wing muscle mechanics in bat flight? *Proc Biol Sci*, 2015. 282(1816): p. 20151832.
32. Hug, F., et al., Nature of the coupling between neural drive and force-generating capacity in the human quadriceps muscle. *Proc Biol Sci*, 2015. 282(1819).
33. Schuermans, J., et al., Biceps femoris and semitendinosus--teammates or competitors? New insights into hamstring injury mechanisms in male football players: a muscle functional MRI study. *Br J Sports Med*, 2014. 48(22): p. 1599-606.